



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

family containing the genera *Balanites*, *Agialida*, and *Agiella* (new); the same author (*idem* 261-271) also breaks up the Boraginaceae into 4 families, Boragaceae, Ehretiaceae, Cordiaceae, and Heliotropiaceae.—C. V. PIPER (Contrib. U. S. Nat. Herb. **10**:1-48. pls. 1-15. 1906), in a revision of N. Am. species of *Festuca*, recognizes 34 and describes 3 as new.—A. A. HELLER (*Muhlenbergia* **2**:177-256. 1906), in an account of botanical exploration in California during 1906, in which numerous species are noted and nomenclatural changes made, describes new species in *Taxiscordion*, *Eriogonum* (2), *Mirabilis*, *Lupinus* (4), *Hesperastragalus*, *Anogra*, *Chylisma*, *Phlox*, *Phacelia*, *Conanthus*, *Cryptanthe* (2), *Amsinckia*, *Pentstemon* (2), and *Orthocarpus*.—PHILIP DOWELL (Bull. Torr. Bot. Club **33**:547-556. pls. 18-22. 1906), in a revision of N. Am. species of *Calceolaria*, recognizes 16 and describes 6 as new.—J. M. C.

**Formative influence of light.**—PEIRCE has extended his studies on irritability in plants from algae to liverworts and ferns, recording the effect of light upon germination and early growth in *Anthoceros fusiformis*, *A. Pearsoni*, *Fimbriaria californica*, and *Gymnogramme triangularis*, and its effect on form in later stages of growth.<sup>23</sup> He finds that germination is dependent upon light of undetermined intensity, and that direction of growth and of successive cell division is determined by the direction of light. The direction of light also profoundly modifies the form of the thalli of the liverworts. This was most marked in *Anthoceros*. On a clinostat they tend to become solid, erect, and cylindrical, conic or vasiform, with radial structure instead of dorsiventral. The author holds that his early results support the hypothesis that, aside from what is actually transmitted from parents to offspring, likeness is due to likeness or identity of the physical environment, and that these factors are as essential determinants as the substance transmitted. Unfortunately PEIRCE'S experiments were interrupted by the earthquake of April 16, so that conclusive tests have not been made in all cases. The work will be continued.—C. R. B.

**Assimilation of organic acids by algae.**—An interesting contribution to our knowledge of the power of assimilation of carbon compounds by autotrophic plants has been made by TRÉBOUX.<sup>24</sup> This investigator experimented with some forty species of the lower algae, testing their power of assimilating various organic acids, which were given in the form of potassium or ammonium salts. Cultures were kept in absolute darkness. It was found that about one-half of the species flourished under the cultural conditions with an organic acid as the only source of carbon; and that of all the acids used acetic acid was most readily assimilated, while acids with larger carbon chains were assimilated in only a few cases. In some cases aminoacids were used with accompanying escape of ammonia. This

<sup>23</sup> PEIRCE, G. J., Studies of irritability in plants. *Annals of Botany* **20**:449-465. pl. 35. 1906.

<sup>24</sup> TRÉBOUX, O., Organische Säuren als Kohlenstoffquelle bei Algen. *Ber. Deutsch. Bot. Gesells.* **23**:432-441. 1906.